



SLOW RELEASE NITROGEN PRODUCTS

THE IMPORTANCE OF SLOW RELEASE NITROGEN

Nitrogen deficiencies can occur throughout the growing season. Symptoms begin on the lower leaves and work upwards. Applying slow release nitrogen (SRN) encourages plant safety and reduces nitrogen leaching. SRN sticks to the foliage for an extended amount of time to supply nitrogen throughout the beginning growth stages. It is also compatible with most herbicides and fungicides acting as a carrier to meet nitrogen needs and treat environmental stresses.



Product & Analysis	Crops	Application Method	Use Rate (per acre)	# of Applications	Estimated Release Pattern	Density (lbs/gal)	Freezing Pt. (°F)
OverPass* CF 22-0-2 + 1.0 S, 0.5 B (25% SRN)	Corn	Foliar	4-8 quarts		10-14 days	10.0	0
	Wheat, barley, other grasses		4-8 pints	1-2			
	Vegetables		2-6 quarts				
OverPass® SF 10-2-10 + 0.5 B, 0.25 Mn (25% SRN)	Soybeans	Foliar	4-8 quarts	2	10-14 days	10.4	24
	Alfalfa/hay		4-8 pints	After each cutting			
	Vegetables		2-6 quarts	Every 14 days when nitrogen is needed			
OverPass® SF Plus 10-2-10 + 0.5 Mn (25% SRN)	Soybeans	Foliar	4-8 quarts	2	10-14 days	10.4	24
	Alfalfa/hay		4-8 pints	After each cutting			
	Vegetables		2-6 quarts	Every 14 days when nitrogen is needed			
Super 72™ 28-0-0 (72% SRN)	Field & row crops; fruits & nuts	Foliar	1-3 gallons	Multiple as needed	25-30 days	10.7	0
Super 25B™ 25-0-0 + 0.5 B (25% SRN)	Field & row crops; fruits & nuts	Foliar	1-2 gallons	Multiple as needed	10-14 days	10.0	0







ROLES OF NITROGEN

- Component of vitamins, amino acids, energy systems
- Increases protein content
- Mobile and moves to root surfaces for absorption



ROLES OF BORON

- Improves seed set under stressful conditions
- · Important cell wall component
- Regulates sugars and starches



ROLES OF SULFUR


- Important in photosynthesis and winter crop hardiness
- Required for synthesis of certain amino acids and proteins



ROLES OF MANGANESE

- Plays vital role in photosynthesis by aiding in chlorophyll synthesis
- Soybeans and wheat require higher amount of Mn

RESEARCH







Summary (Top Left):

A 4.8 bu/A increase was seen in this Wisconsin field test. Season Pass* Plus liquid fertilizer was the starter application used on both treatments. The second treatment received 1 gal/A of OverPass CF liquid fertilizer applied at the V5 stage. This led to a 4.8 bu/A increase over the check.

Summary (Top Right):

The first application of 1 gal/A was made pre-bloom. The second application of 1 gal/A was made at the R1 stage. The combination of the two applications pushed the yield 2.4 bu/A higher than check.

Summary (Bottom): Foliar applied Super 72 resulted in significant yield response, showing a 5-8 bu/A increase over three years in four different locations.



FOR MORE INFORMATION

800-831-4815 png@andersonsinc.com AndersonsPlantNutrient.com



To see how these products can be used in a complete nutrition program visit CROPCOACH.COM